



PROGRAM  
BOOK

XXII Congress

# EuroFoodChem

June 14-16, 2023 | Belgrade, Serbia

<https://xxiieurofoodchem.com>


[congress2023@xxiieurofoodchem.com](mailto:congress2023@xxiieurofoodchem.com)

**Serbian Academy of  
Sciences and Arts**

Knez Mihailova 35  
11000 Belgrade

**Faculty of Chemistry  
University of Belgrade**

Studentski trg 12-16  
11000 Belgrade





# **XXII Congress EuroFoodChem**

**June 14-16, 2023  
Belgrade, Serbia**

*<https://xxiieurofoodchem.com>*

*[congress2023@xxiieurofoodchem.com](mailto:congress2023@xxiieurofoodchem.com)*

## Under the auspices



Ministry of Science, Technological  
Development and Innovations,  
Republic of Serbia



Serbian Academy of  
Sciences and Arts

## Organizers



EuChemS, Division of Food Chemistry



Serbian Chemical Society

- PP 107 **Xiaolu Zhou, Lingjun Ma, Li Dong, Daotong Li, Fang Chen, Xiaosong Hu**  
Bamboo shoot dietary fiber alleviates gut microbiota dysbiosis and modulates liver fatty acid metabolism in mice with high-fat diet-induced obesity
- PP 34 **Natália Čmiková, Simona Kunová, Miroslava Kačániová**  
*Mentha × piperita* L. var. *officinalis* forma *rubescens* Camus antimicrobial activity *in vitro* and *in situ*
- PP 35 **Cristina Martínez-Villaluenga, Irene Tomé-Sánchez, Ana Belén Martín-Diana, Daniel Rico, Iván Jiménez-Pulido, Juana Frias, Vermont P. Dia**  
Anticancer activity of hydrolysed wheat bran mediated through macrophages stimulation
- PP 36 **Bartosz Fotschki, Jerzy Juśkiewicz, Dorota Napórkowska**  
Diet with raspberry polyphenols and prebiotic enhances liver lipid metabolism and regulates synthesis of bile acids in obese rats
- PP 37 **Xiangrong Fang, Yuqing Zhang, Marika Kalpio**  
Health effects and bioavailability of omega-3 fatty acids
- PP 38 **Natalia Platosz, Natalia Bączek, Joanna Topolska, Dorota Szawara-Nowak, Wiesław Wiczowski**  
Chokeberry and red cabbage anthocyanins selectively cross the blood-cerebrospinal fluid barrier
- PP 39 **Lukáš Kolarič, Michaela Lauková, Lucia Minarovičová, Peter Šimko**  
The use of  $\beta$ -cyclodextrin for decreasing the cholesterol content in cereal products
- PP 40 **Ana Sousa, Luis Patarata, Fernando Nunes, Fernanda Cosme**  
Antimicrobial activity of red wine phenolic and non-phenolic fractions
- PP 42 **Ana Cristina Ferrão, Raquel Guiné, Marco Silva, Paula Correia**  
Microbiological and chemical composition of some Portuguese hazelnuts
- PP 43 **Ítala M.G Marx, Elizandra Ardohain, Paula Rodrigues, Andreia Afonso, Tânia C.S.P. Pires, Maria Inês Dias, Lillian Barros, Sandrina A. Heleno**  
Chemical characterization of *Cistus ladanifer* L. phenolic extract and its antifungal activity against *Botrytis cinerea*
- PP 44 **Cristina Caleja, Matilde Rodrigues, André Almeida, Maria Inês Dias, José Pinela, Lillian Barros**  
Fatty acid profile and indices of atherogenicity and thrombogenicity of fish by-product oil for pet food formulations
- PP 45 **Zuzanna Gralak, Kamil Brzuzy, Aneta Jastrzębska**  
Determination of selected parameters and compounds in fermented food

## Microbiological and chemical composition of some Portuguese hazelnuts

Ana Cristina Ferrão<sup>1,2</sup>, Raquel Guiné<sup>1</sup>, Marco Silva<sup>2</sup>, Paula Correia<sup>1</sup>

<sup>1</sup>CERNAS-IPV Research Centre, Polytechnic Institute of Viseu, Viseu Portugal

<sup>2</sup>LAQV/REQUIMTE, NOVA School of Science and Technology, Caparica, Portugal

\* [aferrao@ceex.ipv.pt](mailto:aferrao@ceex.ipv.pt) (Preferred type of presentation: Oral)

Hazelnut is one of the most popular dried fruit all over the world, because of its unique organoleptic characteristics and also due to its nutritional composition [1]. The characteristics of the different hazelnut varieties that exist worldwide are dependent of several factors, such as, the genotype, agricultural and post-harvest practices, climatic conditions and also the geographical location where they are cultivated. Moreover, hazelnut quality is also affected by its chemical composition [2]. The aim of this study was to access the microbiological quality and also the chemical composition of three hazelnut varieties (Grada de Viseu, Tonda de Giffoni and Butler), cultivated in the Viseu region, Portugal. Therefore, hazelnut kernels were analysed for moisture, water activity, fat, protein, fibre and ash. Furthermore, there also quantified the microorganisms at 30°C and the moulds and yeasts at 25°C. All analysis were performed in triplicate. According to the results, fat was the major chemical component for all varieties, varying from 64.38±1.67 g/100 g (var. Grada) to 78.16±1.71 g/100 g (var. Tonda), with statistically significant differences between the varieties ( $p < 0.0005$ ). Moisture content was higher for the var. Butler (6.02±0.37 g/100 g) and lowest for the var. Tonda (4.86±0.33 g/100 g), with statistically significant differences between the three varieties under study ( $p = 0.013$ ). According to the recommendations of the European Union, moisture content of hazelnut kernels should not exceed 6.0% [3]. As for the water activity, the values ranged from 0.54±0.01 (var. Butler) to 0.56±0.01 (var. Grada), but in this case with no statistically significant differences between the varieties. The var. Butler presented the fruits with the higher ash content (2.31±0.18 g/100 g), while var. Grada was the one with the lowest value (1.69±0.16 g/100 g), again with statistically significant differences ( $p = 0.012$ ). The var. Grada presented the higher fibre (6.13±0.03 g/100 g) and protein content (22.84±0.18 g/100 g), with statistically significant differences among the varieties in the two cases. The results also showed (Table 1), that according to the limits established for the count of microorganisms at 30°C and mould and yeast by the National Health Institute Doutor Ricardo Jorge [4] ( $< 10^6$  CFU/g for microorganisms at 30°C,  $< 10^4$  CFU/g for yeast and  $< 5 \times 10^2$  CFU/g for moulds), all the varieties presented a satisfactory microbiological quality, with statistically significant differences between the varieties in the case of the quantification of microorganisms at 30°C. Moreover, as it can be observed in Table 1, var. Butler presented the highest values for the microorganisms at 30°C and also for the moulds and yeast at 25°C, which may be explained by its higher moisture content. The results of this study are very important to characterize the microbiological quality and also some chemical properties of the three most representative hazelnut varieties cultivated in Portugal.

Table 1. Mean count (log CFU/g±standard deviation) of total microorganisms at 30°C and moulds and yeasts at 25°C of the samples under study (n=3).

Sample	Microorganism at 30°C <sup>1</sup>	Mould and Yeasts at 25°C <sup>1</sup>
Grada de Viseu	2.84±0.03 <sup>a</sup>	2.40±0.07
Tonda de Giffoni	2.61±0.03 <sup>a</sup>	2.46±0.04
Butler	2.90±0.01 <sup>a</sup>	2.47±0.03
<i>p</i> -value	<0.005	0.059

<sup>1</sup>Mean values in the same column with the same letter are not statistically different ( $p > 0.05$ )

**Acknowledgments:** We would like to thank to Fundação para a Ciência e Tecnologia (FCT), Portugal, for the Ph.D. scholarship 2021.06088.BD. Thanks also to CERNAS Research Centre and the Polytechnic Institute of Viseu, as well as to LAQV/REQUIMTE of the NOVA School of Science and Technology, for their support. This work was financially supported by Fundação para a Ciência e Tecnologia (FCT), Ph.D. scholarship 2021.06088.BD, with funds from the State Budget and the Community Budget through the European Social Fund of the Union European.

### References:

- [1] X. Wang, E. Tello, D.G. Peterson, *Food Chemistry*, 363 (2021) 130311.
- [2] K. Król, M. Gartner, *Agriculture*, 10 (2020) 375.
- [3] United Nations: Standard DOP-04 concerning the marketing and commercial quality control of Hazelnut Kernels, 2008.
- [4] Instituto Nacional de Saúde Doutor Ricardo Jorge: *Interpretação de resultados de ensaios microbiológicos em alimentos prontos para consumo e em superfícies do ambiente de preparação e distribuição alimentar*. Valores-guia. INSA, 2010.